

Colored Vertex Models, Colored IRF Models and Invariants of Trivalent Colored Graphs

Tetsuo Deguchi and Yasuhiro Akutsu [†]

*Department of Physics, Faculty of Science,
University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan*

[†] *Department of Physics, Faculty of Science
Osaka University, Machikaneyama-cho 1-1, Toyonaka 560, Japan*

Abstract.

We present formulas for the Clebsch-Gordan coefficients and the Racah coefficients for color representations (N -dimensional representations with $q^{2N} = 1$) of $U_q(sl(2))$. We discuss colored vertex models and colored IRF (Interaction Round a Face) models from the color representations of $U_q(sl(2))$. We construct invariants of trivalent colored oriented framed graphs from color representations of $U_q(sl(2))$.

1 Introduction

Recently a new hierarchy of N -state colored braid matrices ($N = 2, 3, \dots$) has been constructed which have color variables α and β . [1, 2, 3, 4] We regard colored braid as a braid on strings with colors. From the colored braid matrices a new series of isotopy invariants of colored links generalizing the multivariable Alexander polynomial has been obtained. [5] Using the limit $q^2 \rightarrow \omega$, we have derived the N -state colored braid matrices from infinite dimensional representations of $U_q(sl(2))$. [6] Here ω is a primitive N -th root of unity. Thus the colored braid matrices are derived from the N -dimensional representations [7] of $U_q(sl(2))$ with $q^{2N} = 1$. We call the N -dimensional representation as (finite dimensional) color representation. We call infinite dimensional representation with a free parameter infinite dimensional color representation. We note that representation theory of quantum groups with q roots of unity is given in Refs. [8, 9].

In this paper we present formulas for the Clebsch-Gordan coefficients and the Racah coefficients of the color representations of $U_q(sl(2))$. We introduce colored IRF models and colored vertex models from the color representations of $U_q(\hat{sl}(2))$. We construct invariants of trivalent colored oriented graphs from the Clebsch-Gordan coefficients of the color representations. The new graph invariants give generalizations of the multivariable Alexander polynomial of links.

There are various viewpoints associated with the colored vertex models. The $N = 2$ case of the colored vertex model corresponds to the trigonometric limit of the Felderhof's solution [10] of the free fermion model. [11] The non-colored case of the trigonometric limit is related to the Lie superalgebra $gl(1|1)$ ($sl(1|1)$). [12] [13, 14, 15] It has been pointed out [16] that the colored case of the trigonometric limit is related to $U_q(gl(1|1))$. From the

different viewpoint it has also been shown [17] that the colored case of the trigonometric limit is related to $U_q(sl(2))$.

The free fermion model and the colored vertex models are related to the link polynomials, which vanish for disconnected links. From the $gl(m|m)$ model we have the Alexander polynomial. [12] A state model for the multi-variable Alexander polynomial was constructed from the trigonometric limit of the free fermion model. [18] Several examples of braid matrices with the vanishing property were calculated by Lee, Couture and Schmeing, by directly solving the braid relation up to the 6 state case. [19]

The outline of this paper is given in the following. In §2 we give the Clebsch-Gordan coefficients and the Racah coefficients of infinite dimensional representations of $U(sl(2))$. In §3 we present formulas for the Clebsch-Gordan coefficients and the Racah coefficients for infinite-dimensional color representations of $U_q(sl(2))$. By taking the limit $q^2 \rightarrow \omega$ we derive formulas for the Clebsch-Gordan coefficients and the Racah coefficients of finite-dimensional color representations of $U_q(sl(2))$. In §4 we discuss the colored vertex models from the viewpoint of color representations of the $U_q(\hat{sl}(2))$. In §5 we construct IRF models associated with color representations of $U_q(sl(2))$, which we have called colored IRF models. In §6 we obtain new invariants of trivalent colored graphs from the Clebsch-Gordan coefficients of color representations.

2 CGC of infinite dimensional representations of $U(sl(2))$

Let us discuss representations of the universal enveloping algebra $U(sl(2))$. The defining relations of the algebra $U(sl(2))$ are given by the following.

$$[H, X^\pm] = \pm 2X^\pm,$$

$$[X^+, X^-] = H. \quad (2.1)$$

The comultiplication is given by

$$\begin{aligned} \Delta(X^\pm) &= X^\pm \otimes I + I \otimes X^\pm, \\ \Delta(H) &= H \otimes I + I \otimes H. \end{aligned} \quad (2.2)$$

Let us define infinite dimensional representation $(\pi^p, V^{(p)})$ of $sl(2)$. Let $V^{(p)}$ be an infinite dimensional vector space over $\mathbf{C}$ with basis $e_0, e_1, \dots$, where e_a is a basis vector in $V^{(p)}$ with the property $(e_a)_b = \delta_{ba}$. We define the matrix elements of the generators as follows.

$$\begin{aligned} (\pi^p(X^+))_b^a &= \sqrt{(2p-a)(a+1)} \cdot \delta_{a+1,b}, \\ (\pi^p(X^-))_b^a &= \sqrt{a(2p-a+1)} \cdot \delta_{a-1,b}, \\ (\pi^p(H))_b^a &= (2p-2a) \cdot \delta_{a,b}. \end{aligned} \quad (2.3)$$

The variable a takes a nonnegative integer; $a = 0, 1, \dots$. The basis vectors for the infinite dimensional representation are labeled by (p, z) , where $p \in \mathbf{C}$ and $z \in \mathbf{Z}_{\geq 0}$. Here the symbol $\mathbf{Z}_{\geq 0}$ denotes the set of nonnegative integers. We write the representation $(\pi^p, V^{(p)})$ simply by $V^{(p)}$, and the basis vector e_z by $|p, z\rangle$. In terms of the basis vector the representation (2.3) is expressed as follows.

$$\begin{aligned} X^+|p, z\rangle &= \sqrt{(2p-z+1)z}|p, z-1\rangle, \\ X^-|p, z\rangle &= \sqrt{(2p-z)(z+1)}|p, z+1\rangle, \\ H|p, z\rangle &= (2p-2z)|p, z\rangle. \end{aligned} \quad (2.4)$$

Let us discuss decomposition of the tensor product $V^{(p_1)} \otimes V^{(p_2)}$. We assume the following

$$|p_1, p_2; p_3, z_3\rangle = \sum_{z_1, z_2} C(p_1, p_2, p_3; z_1, z_2, z_3) |p_1, z_1\rangle \otimes |p_2, z_2\rangle. \quad (2.5)$$

From the comultiplication rule $\Delta(H) = H \otimes I + I \otimes H$ ($H_{tot} = H_1 + H_2$) we can show

$$C(p_1, p_2, p_3; z_1, z_2, z_3) = 0, \text{ unless } p_1 - z_1 + p_2 - z_2 = p_3 - z_3. \quad (2.6)$$

The relation (2.6) gives the charge conservation law. Using (2.6) and the condition $z_i \in \mathbf{Z}_{\geq 0}$, we can show

$$C(p_1, p_2, p_3; z_1, z_2, z_3) = 0, \text{ unless } p_3 = p_1 + p_2 - n, n \in \mathbf{Z}_{\geq 0}. \quad (2.7)$$

Thus we have for $n \in \mathbf{Z}_{\geq 0}$

$$\begin{aligned} & |p_1, p_2; p_1 + p_2 - n, z_3 \rangle \\ &= \sum_{z_1=0}^{z_3+n} C(p_1, p_2, p_1 + p_2 - n; z_1, z_3 + n - z_1, z_3) |p_1, z_1 \rangle \otimes |p_2, z_3 + n - z_1 \rangle. \end{aligned} \quad (2.8)$$

The fusion rules for the tensor product $V^{(p_1)} \otimes V^{(p_2)}$ are summarized as follows.

$$V^{(p_1)} \otimes V^{(p_2)} = \sum_{p_3} N_{p_1 p_2}^{p_3} V^{(p_3)} \quad (2.9)$$

where

$$\begin{aligned} N_{p_1 p_2}^{p_3} &= 1 \text{ for } p_3 = p_1 + p_2 - n, n \in \mathbf{Z}_{\geq 0}, \\ &= 0, \text{ otherwise.} \end{aligned} \quad (2.10)$$

We assume the following convention of the phase factors of the CG coefficients.

$$C(p_1, p_2, p_1 + p_2 - n; 0, n, 0) = 1. \quad (2.11)$$

Then the Clebsch-Gordan coefficients for infinite dimensional representations are given by the following.

$$C(p_1, p_2, p_1 + p_2 - n; z_1, z_2, z_3) = \delta(z_3, z_1 + z_2 - n)$$

$$\begin{aligned}
& \times \sqrt{(2p_1 + 2p_2 - 2n + 1)} \left(\frac{(2p_1 - n)!(2p_2 - n)!(n)!}{(2p_1 + 2p_2 - n + 1)!} \right)^{1/2} \\
& \times \sqrt{(2p_1 - z_1)!(2p_2 - z_2)!(2p_1 + 2p_2 - 2n - z_3)!(z_1)!(z_2)!(z_3)!} \\
& \times \sum_{\nu} \frac{(-1)^{\nu}}{(\nu)!(n - \nu)!} \frac{1}{(z_1 - \nu)!(z_2 - n + \nu)!} \\
& \times \frac{1}{(2p_1 - n - z_1 + \nu)!(2p_2 - z_2 - \nu)!}.
\end{aligned} \tag{2.12}$$

Here we have defined the symbol $(p)!$ through the Gamma function

$$(p)! = \Gamma(p + 1). \tag{2.13}$$

The sum over the integer ν in (2.12) is taken under the following condition

$$\max\{0, n - z_2\} \leq \nu \leq \min\{n, z_1\}. \tag{2.14}$$

We can derive the Clebsch-Gordan coefficients (2.12) following Racah's derivation [20] (see also Appendix A).

Hereafter we assume that the symbols p and p_j denote complex parameters ($p, p_j \in \mathbf{C}$), and $2j$ and $2j_k$ denote nonnegative integers ($2j, 2j_k \in \mathbf{Z}_{\geq 0}$). We also assume that $2\mu_j$ ($j = 1, 2, 3$) denote either non-negative integers ($2\mu_j \in \mathbf{Z}_{\geq 0}$) or complex parameters ($\mu_j \in \mathbf{C}$).

Let $V^{(j)}$ denote the representation with dimension $2j + 1$ (spin j representation). Let us consider the following decomposition of the tensor product.

$$V^{(\mu_1)} \otimes V^{(\mu_2)} = \sum_{p_3} N_{\mu_1 \mu_2}^{p_3} V^{(p_3)}, p_3 \in \mathbf{C}. \tag{2.15}$$

Here we consider the two cases, (1) $(\mu_1, \mu_2) = (p_1, j_2)$ ($p_1 \in \mathbf{C}$, $2j_2 \in \mathbf{Z}_{\geq 0}$); (2) $(\mu_1, \mu_2) = (j_1, p_2)$ ($2j_1 \in \mathbf{Z}_{\geq 0}$, $p_2 \in \mathbf{C}$).

Then we have the following fusion rules.

Case (1) $(\mu_1, \mu_2) = (p_1, j_2)$ ($p_1 \in \mathbf{C}$, $2j_2 \in \mathbf{Z}_{\geq 0}$);

$$\begin{aligned}
N_{p_1 j_2}^{p_3} &= 1 \text{ if } p_3 = p_1 + j_2 - n, 0 \leq n \leq 2j_2, n \in \mathbf{Z}_{\geq 0}, \\
&= 0, \quad \text{otherwise.}
\end{aligned} \tag{2.16}$$

Case (2) $(\mu_1, \mu_2) = (j_1, p_2)$ ($2j_1 \in \mathbf{Z}_{\geq 0}$, $p_2 \in \mathbf{C}$)

$$\begin{aligned} N_{j_1 p_2}^{p_3} &= 1 \text{ if } p_3 = j_1 + p_2 - n, 0 \leq n \leq 2j_1, n \in \mathbf{Z}_{\geq 0}, \\ &= 0, \quad \text{otherwise.} \end{aligned} \quad (2.17)$$

The Clebsch-Gordan coefficients are given by the following.

$$\begin{aligned} C(\mu_1, \mu_2, \mu_1 + \mu_2 - n; z_1, z_2, z_3) &= \delta(z_3, z_1 + z_2 - n) \\ &\times \sqrt{(2\mu_1 + 2\mu_2 - 2n + 1)} \left(\frac{(\mu_1 - n)!(2\mu_2 - n)!(n)!}{(2\mu_1 + 2\mu_2 - n + 1)!} \right)^{1/2} \\ &\times \sqrt{(2\mu_1 - z_1)!(2\mu_2 - z_2)!(2\mu_1 + 2\mu_2 - 2n - z_3)!(z_1)!(z_2)!(z_3)!} \\ &\times \sum_{\nu} \frac{(-1)^{\nu}}{(\nu)!(n - \nu)!} \frac{1}{(z_1 - \nu)!(z_2 - n + \nu)!} \\ &\times \frac{1}{(2\mu_1 - n - z_1 + \nu)!(2\mu_2 - z_2 - \nu)!} \end{aligned} \quad (2.18)$$

Here the sum over the integer ν is taken under the following conditions:

Case (1) $\mu_1 = p_1, \mu_2 = j_2$,

$$\max\{0, n - z_2\} \leq \nu \leq \min\{n, z_1, 2j_2 - z_2\}. \quad (2.19)$$

Case (2) $\mu_1 = j_1, \mu_2 = p_2$,

$$\max\{0, n - z_2, n + z_1 - 2j_1\} \leq \nu \leq \min\{n, z_1\}. \quad (2.20)$$

We note that the formula (2.18) can be derived from (2.12) by taking the limit $p_2 \rightarrow j_2$. The branching rule (2.17) can be shown through this limiting procedure.

The Clebsch-Gordan coefficients satisfy the orthogonality relations for the cases $(\mu_1, \mu_2) = (p_1, p_2)$, (p_1, j_2) , and (j_1, p_2) .

$$\begin{aligned} \delta_{nn'} &= \sum_{z_1} C(\mu_1, \mu_2, \mu_1 + \mu_2 - n; z_1, \tau - z_1, \tau - n) \\ &\quad \times C(\mu_1, \mu_2, \mu_1 + \mu_2 - n'; z_1, \tau - z_1, \tau - n'). \end{aligned} \quad (2.21)$$

Let us discuss the Racah coefficients for finite and infinite dimensional representations. Using the formulas for the Clebsch-Gordan coefficients (2.12), and (2.18) we can calculate the Racah coefficients. We recall that $2\mu_i$ ($i = 1, 2, 3$) is either a nonnegative integer ($2\mu_i \in \mathbf{Z}_{\geq 0}$) or a complex parameter ($\mu_i \in \mathbf{C}$). We introduce μ, μ', μ'' by

$$\mu = \mu_1 + \mu_2 + \mu_3 - n, \mu' = \mu_1 + \mu_2 - n', \mu'' = \mu_2 + \mu_3 - n'', \quad (2.22)$$

where $n, n', n'' \in \mathbf{Z}_{\geq 0}$.

The Racah coefficients $W(\mu_1, \mu_2, \mu, \mu_3; \mu', \mu'')$ are defined by [20]

$$\begin{aligned} & |\mu_1 \mu_2 (\mu') \mu_3 \mu z \rangle = \\ & = \sum_{\mu''} \sqrt{(2\mu' + 1)(2\mu'' + 1)} W(\mu_1, \mu_2, \mu, \mu_3; \mu', \mu'') |\mu_1, \mu_2 \mu_3 (\mu'') \mu z \rangle. \end{aligned} \quad (2.23)$$

We note that the Racah coefficients do not depend on z by the definition (2.23). From the definition of the Racah coefficients, we have

$$\begin{aligned} & C(\mu_1, \mu_2, \mu'; w_1, w_2, w_1 + w_2 - n') C(\mu', \mu_3, \mu; w_1 + w_2 - n', w_3, w_1 + w_2 + w_3 - n) \\ & = \sum_{\mu''} R_{\mu'' \mu'} C(\mu_1, \mu'', \mu; w_1, w_2 + w_3 - n'', w_1 + w_2 + w_3 - n) \\ & \quad \times C(\mu_2, \mu_3, \mu''; w_2, w_3, w_2 + w_3 - n''). \end{aligned} \quad (2.24)$$

Here $w_1, w_2, w_3 \in \mathbf{Z}_{\geq 0}$. Applying the orthogonality relation (2.21) to (2.24) we have a formula for the Racah coefficients. The derivation of the Racah coefficients consists of the following procedures. (1) Let us set $w_2 + w_3 = z''$. (2) Then multiply both the left hand side and right hand side of (2.24) by $\sum_{w_2} C(\mu_2, \mu_3, \mu_2 + \mu_3 - N''; w_2, z'' - w_2, z'' - N'')$. (3) Rewrite N'' by n'' , and introduce $z = w_1 + z'' = w_1 + w_2 + w_3$. (4) Multiply both sides by $\sum_{w_1} C(\mu_1, \mu_2 + \mu_3 - n'', \mu_1 + \mu_2 + \mu_3 - n; w_1, z - w_1, z + n'' - n)$.

Thus we have the Racah coefficients $W(\mu_1, \mu_2, \mu, \mu_3; \mu', \mu'')$ as follows.

$$\begin{aligned}
R_{\mu'' \mu'} &= \sqrt{(2\mu' + 1)(2\mu'' + 1)} W(\mu_1, \mu_2, \mu, \mu_3; \mu', \mu'') \\
&= \sum_{w_1} \sum_{w_2} C(\mu_1, \mu_2, \mu'; w_1, w_2, w_1 + w_2 - n') \\
&\quad \times C(\mu', \mu_3, \mu; w_1 + w_2 - n', z - w_1 - w_2, z - n) \\
&\quad \times C(\mu_2, \mu_3, \mu_2 + \mu_3 - n''; w_2, z - w_1 - w_2, z - w_1 - n'') \\
&\quad \times C(\mu_1, \mu'', \mu; w_1, z - w_1, z + n'' - n). \tag{2.25}
\end{aligned}$$

3 CGC of color representations of $U_q(sl(2))$

3.1 Infinite dimensional case

The generators of the algebra are $\{X^+, X^-, H\}$ with the defining relations [21, 22]

$$\begin{aligned}
[H, X^\pm] &= \pm 2X^\pm, \\
[X^+, X^-] &= \frac{q^H - q^{-H}}{q - q^{-1}}. \tag{3.1}
\end{aligned}$$

The comultiplication is given by

$$\begin{aligned}
\Delta(H) &= H \otimes I + I \otimes H, \\
\Delta(X^\pm) &= X^\pm \otimes q^{H/2} + q^{-H/2} \otimes X^\pm. \tag{3.2}
\end{aligned}$$

We use the following symbols for q -analogs.

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}, \quad [n]_q! = \prod_{k=1}^n [k]_q, \quad [p; n]_q! = \prod_{k=0}^{n-1} [p - k]_q. \tag{3.3}$$

Here n is a positive integer. For the case $n = 0$ we assume

$$[0]_q! = [p; 0]_q! = 1. \tag{3.4}$$

Let us define an infinite dimensional representation $(\pi_q^p, V^{(p)})$ of $U_q(sl(2))$. We recall that p is a complex parameter. Let $V^{(p)}$ be an infinite dimensional vector space over $\mathbf{C}$ with basis $e_0, e_1, \dots$, where e_a is a basis vector in $V^{(p)}$ with the property $(e_a)_b = \delta_{ba}$. We define matrix elements of the representations of the generators X^+ , X^- and $K = q^H$ for $(\pi_q^p, V^{(p)})$ as follows.

$$\begin{aligned} (\pi^p(X^+)_q)_b^a &= \sqrt{[2p-a]_q[a+1]_q} \cdot \delta_{a+1,b}, \\ (\pi^p(X^-)_q)_b^a &= \sqrt{[2p-a+1]_q[a]_q} \delta_{a-1,b}, \\ (\pi^p(H)_q)_b^a &= (2p-2a) \cdot \delta_{a,b}. \end{aligned} \quad (3.5)$$

Here a, b are nonnegative integers ($a, b = 0, 1, \dots$), and $p \in \mathbf{C}$. It is easy to see that the operators defined in (3.5) satisfy the defining relations of the algebra $U_q(sl(2))$ with q generic. We call the representation in (3.5) infinite dimensional color representation. We write the basis vector $e_a(p)$ by $|p, z >_q$, and the representation $(\pi_q^p, V^{(p)})$ by $V_q^{(p)}$.

Let us discuss decomposition of the tensor product $V_q^{p_1} \otimes V_q^{(p_2)}$. We define the Clebsch-Gordan coefficients by the following

$$|p_1, p_2; p_3, z_3 >_q = \sum_{z_1, z_2} C(p_1, p_2, p_3; z_1, z_2, z_3)_q |p_1, z_1 >_q \otimes |p_2, z_2 >_q. \quad (3.6)$$

From the comultiplication rule we can show

$$C(p_1, p_2, p_3; z_1, z_2, z_3)_q = 0, \text{ unless } p_1 - z_1 + p_2 - z_2 = p_3 - z_3. \quad (3.7)$$

The relation (3.7) leads to the charge conservation law. Using (3.7) and the condition $z_i \in \mathbf{Z}_{\geq 0}$, we can show

$$C(p_1, p_2, p_3; z_1, z_2, z_3)_q = 0, \text{ unless } p_3 = p_1 + p_2 - n, n \in \mathbf{Z}_{\geq 0}. \quad (3.8)$$

Thus we have following.

$$|p_1, p_2; p_1 + p_2 - n, z_3 >_q$$

$$= \sum_{z_1=0}^{z_3+n} C(p_1, p_2, p_1 + p_2 - n; z_1, z_3 + n - z_1, z_3)_q |p_1, z_1 \rangle_q \otimes |p_2, z_3 + n - z_1 \rangle_q . \quad (3.9)$$

Here $n \in \mathbf{Z}_{\geq 0}$. The Clebsch-Gordan coefficients for infinite dimensional representations are given by the following.

$$\begin{aligned} C(p_1, p_2, p_1 + p_2 - n; z_1, z_2, z_3)_q &= \delta(z_3, z_1 + z_2 - n) \\ &\times \sqrt{[2p_1 + 2p_2 - 2n + 1]_q} \sqrt{[n]_q! [z_1]_q! [z_2]_q! [z_3]_q!} \\ &\times q^{(n-n^2)/2 + (n-z_2)p_1 + (n+z_1)p_2} \\ &\times \sum_{\nu} \frac{(-1)^{\nu} q^{-\nu(2p_1+2p_2-n+1)}}{[\nu]_q! [n-\nu]_q! [z_1-\nu]_q! [z_2-n+\nu]_q!} \\ &\times \sqrt{\frac{[2p_1-n; z_1-\nu]_q! [2p_1-z_1; n-\nu]_q! [2p_2-n; z_2+\nu-n]_q! [2p_2-z_2; \nu]_q!}{[2p_1+2p_2-n+1; z_1+z_2+1]_q!}}. \end{aligned} \quad (3.10)$$

Here $p_3 = p_1 + p_2 - n, n \in \mathbf{Z}_{\geq 0}$, and the sum over the integer ν in (3.10) is taken under the following condition

$$\max\{0, n - z_2\} \leq \nu \leq \min\{n, z_1\}. \quad (3.11)$$

We shall derive the symmetric expression (3.10) of the Clebsch-Gordan coefficients (3.10) in Appendix A.

Let $V_q^{(j)}$ be the spin j representation of $U_q(sl(2))$. Let us discuss decomposition of the tensor product $V_q^{(p_1)} \otimes V_q^{(j_2)}$ and $V_q^{(j_1)} \otimes V_q^{(p_2)}$. It is easy to see the following rules.

$$\begin{aligned} C(\mu_1, \mu_2, p_3; z_1, z_2, z_3)_q &= 0, \\ \text{unless } \mu_1 - z_1 + \mu_2 - z_2 &= p_3 - z_3, p_3 = \mu_1 + \mu_2 - n \quad (n \in \mathbf{Z}_{\geq 0}), \\ \text{and } \begin{cases} 0 \leq n \leq 2j_2, 0 \leq z_2 \leq 2j_2, & \text{if } (\mu_1, \mu_2) = (p_1, j_2), \\ 0 \leq n \leq 2j_1, 0 \leq z_1 \leq 2j_1, & \text{if } (\mu_1, \mu_2) = (j_1, p_2). \end{cases} \end{aligned} \quad (3.12)$$

The Clebsch-Gordan coefficients are given by

$$\begin{aligned}
C(\mu_1, \mu_2, \mu_1 + \mu_2 - n; z_1, z_2, z_3)_q &= \delta(z_3, z_1 + z_2 - n) \\
&\times \sqrt{[2\mu_1 + 2\mu_2 - 2n + 1]_q} \sqrt{[n]_q! [z_1]_q! [z_2]_q! [z_3]_q!} \\
&\times q^{(n-n^2)/2 + (n-z_2)\mu_1 + (n+z_1)\mu_2} \\
&\times \sum_{\nu} \frac{(-1)^{\nu} q^{-\nu(2\mu_1+2\mu_2-n+1)}}{[\nu]_q! [n-\nu]_q! [z_1-\nu]_q! [z_2-n+\nu]_q!} \\
&\times \left(\frac{[2\mu_1-n; z_1-\nu]_q! [2\mu_1-z_1; n-\nu]_q! [2\mu_2-n; z_2+\nu-n]_q! [2\mu_2-z_2; \nu]_q!}{[2\mu_1+2\mu_2-n+1; z_1+z_2+1]_q!} \right)^{1/2}.
\end{aligned} \tag{3.13}$$

Here the sum over the integer ν in (3.13) is taken under the following conditions.

Case (1): $\mu_1 = p_1, \mu_2 = j_2$

$$\max\{0, n - z_2\} \leq \nu \leq \min\{n, z_1, 2j_2 - z_2\}. \tag{3.14}$$

Case (2): $\mu_1 = j_1, \mu_2 = p_2$

$$\max\{0, n - z_2, n + z_1 - 2j_1\} \leq \nu \leq \min\{n, z_1\}. \tag{3.15}$$

For convenience we give the Clebsch-Gordan coefficients for finite dimensional representations $V^{(j_i)}$, ($i = 1, 2, 3$) in the following [23] (see also [24]).

$$\begin{aligned}
C(j_1, j_2, j_3; z_1, z_2, z_3)_q &= \delta(z_3, z_1 + z_2 - n) \\
&\times ([2j_3 + 1]_q)^{1/2} \Delta_q(j_1 j_2 j_3) \\
&\times q^{j_1(j_1+1) + j_2(j_2+1) - j_3(j_3+1) + 2(j_1 j_2 + j_1(j_2 - z_2) - j_2(j_1 - z_1)) / 2} \\
&\times ([2j_1 - z_1]_q! [z_1]_q! [2j_2 - z_2]_q! [z_2]_q! [2j_3 - z_3]_q! [z_3]_q!)^{1/2} \\
&\times \sum_{\nu} \frac{(-1)^{\nu} q^{-\nu(j_1+j_2+j_3+1)}}{[\nu]_q! [j_1+j_2-j_3-\nu]_q! [z_1-\nu]_q!} \\
&\times \frac{1}{[2j_2-z_2-\nu]_q! [j_3+j_1-j_2-z_1+\nu]_q! [j_3-j_1-j_2+z_2+\nu]_q!}.
\end{aligned} \tag{3.16}$$

Here the sum over the integer ν is such that

$$\max\{0, j_2 - j_3 - j_1 + z_1, j_1 + j_2 - j_3 - z_2\} \leq \nu \leq \min\{z_1, j_1 + j_2 - j_3, 2j_2 - z_2\}. \quad (3.17)$$

The symbol $\Delta_q(abc)$ has been defined by

$$\Delta_q(abc) = \left(\frac{[a+b-c]_q! [c+a-b]_q! [b+c-a]_q!}{[a+b+c+1]_q!} \right)^{1/2}. \quad (3.18)$$

If we replace p_i by j_i ($i = 1, 2, 3$), then the formal expression of the Clebsch-Gordan coefficients for infinite dimensional representations is consistent with the finite dimensional one (3.16).

The Clebsch-Gordan coefficients satisfy the orthogonality relations for the cases $(\mu_1, \mu_2) = (p_1, p_2)$, (p_1, j_2) , and (j_1, p_2) .

$$\begin{aligned} \delta_{nn'} &= \sum_{z_1} C(\mu_1, \mu_2, \mu_1 + \mu_2 - n; z_1, \tau - z_1, \tau - n)_q \\ &\quad \times C(\mu_1, \mu_2, \mu_1 + \mu_2 - n'; z_1, \tau - z_1, \tau - n')_q, \end{aligned} \quad (3.19)$$

We can calculate the Racah coefficients using the Clebsch-Gordan coefficients and the orthogonality relations (3.19) as in §2. The Racah coefficients are given by

$$\begin{aligned} R_{\mu''\mu'}(q) &= \sqrt{[2\mu''+1]_q [2\mu'+1]_q} W(\mu_1, \mu_2, \mu, \mu_3; \mu', \mu'')_q \\ &= \sum_{w_1} \sum_{w_2} C(\mu_1, \mu_2, \mu'; w_1, w_2, w_1 + w_2 - n')_q \\ &\quad \times C(\mu', \mu_3, \mu; w_1 + w_2 - n', z'' - w_2, w_1 + z'' - n)_q \\ &\quad \times C(\mu_2, \mu_3, \mu_2 + \mu_3 - n''; w_2, z'' - w_2, z'' - n'')_q \\ &\quad \times C(\mu_1, \mu'', \mu; w_1, z'', w_1 + z'' + n'' - n)_q. \end{aligned} \quad (3.20)$$

Here we have assumed that $2\mu_i$ ($i = 1, 2, 3$) is either a nonnegative integer ($2\mu_i \in \mathbf{Z}_{\geq 0}$) or a complex parameter ($\mu_i \in \mathbf{C}$). We have defined μ, μ', μ'' by

$$\mu = \mu_1 + \mu_2 + \mu_3 - n, \mu' = \mu_1 + \mu_2 - n', \mu'' = \mu_2 + \mu_3 - n''. \quad (3.21)$$

where $n, n', n'' \in \mathbf{Z}_{\geq 0}$.

3.2 Finite dimensional cases

Let ω be a primitive N -th root of unity:

$$\omega = \exp\left(\frac{2\pi i s}{N}\right), \quad (N, s) = 1. \quad (3.22)$$

Here the symbol $(a, b) = 1$ denotes that the integers a and b have no common divisor except 1. We write by ϵ a square root of ω : $\epsilon = \omega^{1/2} = \exp(\pi i s/N)$, $(N, s) = 1$.

We take the limit $q \rightarrow \epsilon$ in the infinite dimensional representation (3.5). Then we have

$$\lim_{q \rightarrow \epsilon} \left(\pi^p(X^-)_q \right)_{N-1}^N = 0. \quad (3.23)$$

From the property (3.23) we can restrict the infinite dimensions into N dimensions: $V_q^{(p)}(\infty) \rightarrow V_\epsilon^{(p)}(N)$, where $V_\epsilon^{(p)}(N)$ is an N -dimensional vector space with basis $e_0, \dots, e_{N-1}$. For $a, b = 0, 1, \dots, N-1$ we have the following matrix representations

$$\begin{aligned} \left(\pi^p(X^+)_{q=\epsilon} \right)_b^a &= ([2p-a]_\epsilon [a+1]_\epsilon)^{1/2} \cdot \delta_{a+1,b}, \\ \left(\pi^p(X^-)_{q=\epsilon} \right)_b^a &= ([2p-a+1]_\epsilon [a]_\epsilon)^{1/2} \cdot \delta_{a-1,b}, \\ \left(\pi^p(K)_{q=\epsilon} \right)_{ab} &= \epsilon^{(2p-2a)} \cdot \delta_{a,b}. \end{aligned} \quad (3.24)$$

We write the finite dimensional color representation $\{\pi_\epsilon^p, V_\epsilon^{(p)}\}$ by $V_\epsilon^{(p)}$. We write the basis vector e_a by $|p, z \rangle_\epsilon$. The basis vectors for $V_\epsilon^{(p)}$ are $\{|p, z \rangle_\epsilon, z = 0, \dots, N-1\}$.

Let us discuss decomposition of the tensor product $V_\epsilon^{(p_1)} \otimes V_\epsilon^{(p_2)}$. We take the limit $q \rightarrow \epsilon$ in the expression of the Clebsch-Gordan coefficients (3.10). Since there is no singularity in the limiting process in the expression of the

Clebsch-Gordan coefficients, we have the fusion rule in the following.

$$V_\epsilon^{(p_1)} \otimes V_\epsilon^{(p_2)} = \sum_{p_3} N_{p_1, p_2}^{p_3} V_\epsilon^{(p_3)}, \quad (3.25)$$

where

$$\begin{aligned} N_{p_1, p_2}^{p_3} &= 1, \text{ for } p_3 = p_1 + p_2, p_1 + p_2 - 1, \dots, p_1 + p_2 - N + 1. \\ &= 0, \text{ otherwise.} \end{aligned} \quad (3.26)$$

We note that the condition $0 \leq n \leq N - 1$ is derived from the factor $\sqrt{[n]_q!}$ in (3.10). In terms of the Clebsch-Gordan coefficients we have

$$\begin{aligned} C(p_1, p_2, p_3; z_1, z_2, z_3)_\epsilon &= 0, \\ \text{unless } p_1 - z_1 + p_2 - z_2 &= p_3 - z_3, p_3 = p_1 + p_2 - n, \\ 0 \leq z_i \leq N - 1 \text{ for } i = 1, 2, 3, \text{ and } 0 \leq n \leq N - 1, n \in \mathbf{Z}_{\geq 0}. \end{aligned} \quad (3.27)$$

By taking the limit $q \rightarrow \epsilon$ in (3.10) we obtain the Clebsch-Gordan coefficients for finite dimensional color representations. Let us set $p_3 = p_1 + p_2 - n$, where n is an integer with $0 \leq n \leq N - 1$, and $0 \leq z_i \leq N - 1$, for $i=1,2,3$. The Clebsch-Gordan coefficients for $V_\epsilon^{(p_i)}$ for $i = 1, 2, 3$ are given as follows.

$$\begin{aligned} C(p_1, p_2, p_1 + p_2 - n; z_1, z_2, z_3)_\epsilon &= \delta(z_3, z_1 + z_2 - n) \\ &\times \sqrt{[2p_1 + 2p_2 - 2n + 1]_\epsilon} \sqrt{[n]_\epsilon! [z_1]_\epsilon! [z_2]_\epsilon! [z_3]_\epsilon!} \\ &\times \epsilon^{(n-n^2)/2 + (n-z_2)p_1 + (n+z_1)p_2} \\ &\times \sum_\nu \frac{(-1)^\nu \epsilon^{-\nu(p_1 + p_2 + p_3 + 1)}}{[\nu]_\epsilon! [n - \nu]_\epsilon! [z_1 - \nu]_\epsilon! [z_2 - n + \nu]_\epsilon!} \\ &\times \sqrt{\frac{[2p_1 - n; z_1 - \nu]_\epsilon! [2p_1 - z_1; n - \nu]_\epsilon! [2p_2 - n; z_2 + \nu - n]_\epsilon! [2p_2 - z_2; \nu]_\epsilon!}{[2p_1 + 2p_2 - n + 1; z_1 + z_2 + 1]_\epsilon!}}. \end{aligned} \quad (3.28)$$

Here the sum over the integer ν in (3.28) is taken under the following condition

$$\max\{0, n - z_2\} \leq \nu \leq \min\{n, z_1\}. \quad (3.29)$$

Let $V_\epsilon^{(j)}$ be the spin j representation of $U_\epsilon(sl(2))$. When q is a root of unity ($q^{2N} = 1$), the spin j representations of $U_q(sl(2))$ has the following constraint. [25, 7]

$$0 \leq 2j \leq N - 2, \quad 2j \in \mathbf{Z}_{\geq 0}. \quad (3.30)$$

Let us discuss decomposition of the tensor product

$$V_\epsilon^{(\mu_1)} \otimes V_\epsilon^{(\mu_2)} = \sum_{p_3} N_{\mu_1 \mu_2}^{p_3} V_\epsilon^{(p_3)} \quad (3.31)$$

where $(\mu_1, \mu_2) = (p_1, j_2), (j_1, p_2)$. By taking the limit $q \rightarrow \epsilon$ in (3.13) we have the following fusion rule.

Case (1): $\mu_1 = p_1, \mu_2 = j_2 \quad (2j_2 < N - 1)$

$$\begin{aligned} N_{p_1 j_2}^{p_3} &= 1 \text{ for } p_3 = p_1 + j_2 - n, \quad 0 \leq n \leq 2j_2, n \in \mathbf{Z}, \\ &= 0, \quad \text{otherwise.} \end{aligned} \quad (3.32)$$

Case (2): $\mu_1 = j_1, \mu_2 = p_2 \quad (2j_1 < N - 1)$

$$\begin{aligned} N_{j_1 p_2}^{p_3} &= 1 \text{ for } p_3 = j_1 + p_2 - n, \quad 0 \leq n \leq 2j_1, n \in \mathbf{Z}, \\ &= 0, \quad \text{otherwise.} \end{aligned} \quad (3.33)$$

By taking the limit: $q \rightarrow \epsilon$ in (3.13), we have the Clebsch-Gordan coefficients. Let us set $p_3 = \mu_1 + \mu_2 - n$. When $(\mu_1, \mu_2) = (p_1, j_2)$, we assume that $0 \leq z_i \leq N - 1$ ($i=1,3$), $0 \leq z_2 \leq 2j_2$ and $0 \leq n \leq 2j_2$. When $(\mu_1, \mu_2) = (j_1, p_2)$, we assume that $0 \leq z_i \leq N - 1$ ($i=2,3$), $0 \leq z_1 \leq 2j_1$ and $0 \leq n \leq 2j_1$. The Clebsch-Gordan coefficients are given in the following.

$$C(\mu_1, \mu_2, \mu_1 + \mu_2 - n; z_1, z_2, z_3)_\epsilon = \delta(z_3, z_1 + z_2 - n)$$

$$\begin{aligned}
& \times \sqrt{[2\mu_1 + 2\mu_2 - 2n + 1]_\epsilon} \sqrt{[n]_\epsilon! [z_1]_\epsilon! [z_2]_\epsilon! [z_3]_\epsilon!} \\
& \times \epsilon^{(n-n^2)/2 + (n-z_2)\mu_1 + (n+z_1)\mu_2} \\
& \times \sum_{\nu} \frac{(-1)^\nu \epsilon^{-\nu(2\mu_1+2\mu_2-n+1)}}{[\nu]_\epsilon! [n-\nu]_\epsilon! [z_1-\nu]_\epsilon! [z_2-n+\nu]_\epsilon!} \\
& \times \left(\frac{[2\mu_1-n; z_1-\nu]_\epsilon! [2\mu_1-z_1; n-\nu]_\epsilon! [2\mu_2-n; z_2+\nu-n]_\epsilon! [2\mu_2-z_2; \nu]_\epsilon!}{[2\mu_1+2\mu_2-n+1; z_1+z_2+1]_\epsilon!} \right)^{1/2}.
\end{aligned} \tag{3.34}$$

Here the sum over the integer ν in (3.34) is taken under the following condition.

Case (1): $\mu_1 = p_1, \mu_2 = j_2$

$$\max\{0, n - z_2\} \leq \nu \leq \min\{n, z_1, 2j_2 - z_2\}. \tag{3.35}$$

Case (2): $\mu_1 = j_1, \mu_2 = p_2$

$$\max\{0, n - z_2, n + z_1 - 2j_1\} \leq \nu \leq \min\{n, z_1\}. \tag{3.36}$$

The Clebsch-Gordan coefficients have the following orthogonality relations for the cases $(\mu_1, \mu_2) = (p_1, p_2), (p_1, j_2), (j_1, p_2)$.

$$\begin{aligned}
\delta_{nn'} &= \sum_{z_1} C(\mu_1, \mu_2, \mu_1 + \mu_2 - n; z_1, \tau - z_1, \tau - n)_\epsilon \\
&\quad \times C(\mu_1, \mu_2, \mu_1 + \mu_2 - n'; z_1, \tau - z_1, \tau - n')_\epsilon,
\end{aligned} \tag{3.37}$$

$$\begin{aligned}
\delta_{z_1 z'_1} &= \sum_n C(\mu_1, \mu_2, \mu_1 + \mu_2 - n; z_1, z + n - z_1, z)_\epsilon \\
&\quad \times C(\mu_1, \mu_2, \mu_1 + \mu_2 - n; z'_1, z + n - z'_1, z)_\epsilon.
\end{aligned} \tag{3.38}$$

Recall that $2\mu_i$ ($i = 1, 2, 3$) be either a nonnegative integer ($2\mu_i \in \mathbf{Z}_{\geq 0}$) or a complex parameter ($\mu_i \in \mathbf{C}$). We introduce μ, μ', μ'' by

$$\mu = \mu_1 + \mu_2 + \mu_3 - n, \mu' = \mu_1 + \mu_2 - n', \mu'' = \mu_2 + \mu_3 - n''. \tag{3.39}$$

where $n, n', n'' \in \mathbf{Z}_{\geq 0}$. In terms of the Clebsch-Gordan coefficients the Racah coefficients $W(\mu_1, \mu_2, \mu, \mu_3; \mu', \mu'')_\epsilon$ are given by the following.

$$\begin{aligned}
R_{\mu'' \mu'}(\epsilon) &= \sqrt{[2\mu' + 1]_\epsilon [2\mu'' + 1]_\epsilon} W(\mu_1, \mu_2, \mu, \mu_3; \mu', \mu'')_\epsilon \\
&= \sum_{w_1} \sum_{w_2} C(\mu_1, \mu_2, \mu'; w_1, w_2, w_1 + w_2 - n')_\epsilon \\
&\quad \times C(\mu', \mu_3, \mu; w_1 + w_2 - n', z'' - w_2, w_1 + z'' - n)_\epsilon \\
&\quad \times C(\mu_2, \mu_3, \mu_2 + \mu_3 - n''; w_2, z'' - w_2, z'' - n'')_\epsilon \\
&\quad \times C(\mu_1, \mu'', \mu; w_1, z'', w_1 + z'' + n'' - n)_\epsilon. \tag{3.40}
\end{aligned}$$

Finally we give comments. (1) The finite dimensional color representation in this section is equivalent to the N dimensional representation of $U_q(sl(2))$ with $q^{2N} = 1$ in the reference [7]. We have discussed the representation from the view point of the infinite dimensional representation of $U_q(sl(2))$ and the limit $q \rightarrow \epsilon$ introduced in the reference [6], and then through the limit we have obtained the formulas for the Clebsch-Gordan coefficients and the Racah coefficients. (2) Fusion rules similar to (3.26) and (3.32) have been given in the reference [26]. The fusion rules are for the m dimensional representations (semi-periodic representations) of $U_q(sl(2))$ with $q^m = 1$, which are different from the finite dimensional color representations (the N dimensional representation with $q^{2N} = 1$).

4 Colored vertex models

4.1 Quantum affine algebra $U_q(\hat{sl}(2))$

We express the Boltzmann weights of the colored vertex models in terms of the Clebsch-Gordan coefficients of color representations. Let us consider the q -analog of the universal enveloping algebra $U_q(\hat{sl}(2, C))$ of the affine

Kac-Moody algebra $\hat{sl}(2)$. [22] The generators $\{X_i^\pm, H_i; i = 0, 1\}$ satisfy the following defining relations $(i, j = 0, 1)$.

$$\begin{aligned}
[H_i, H_j] &= 0, \\
[H_i, X_i^\pm] &= \pm 2X_i^\pm, \quad [H_i, X_j^\pm] = \mp 2X_j^\pm \quad (i \neq j), \\
[X_i^+, X_j^-] &= \delta_{ij} \frac{q^{H_i} - q^{-H_i}}{q - q^{-1}}, \\
\sum_{\nu=0}^3 (-1)^\nu \frac{[3]_q!}{[3-\nu]_q! [\nu]_q!} (X^\pm)^{3-\nu} X_j^\pm (X_i^\pm)^\nu &= 0 \quad (i \neq j). \quad (4.1)
\end{aligned}$$

The comultiplication is given by

$$\begin{aligned}
\Delta(H_i) &= H_i \otimes I + I \otimes H_i, \\
\Delta(X_i^\pm) &= X_i^\pm \otimes q^{H_i/2} + q^{-H_i/2} \otimes X_i^\pm. \quad (4.2)
\end{aligned}$$

We denote by $\mathcal{R}$ the universal R matrix of $U_q(\hat{sl}(2, C))$. The universal R matrix satisfies the following

$$\mathcal{R}\Delta(a) = \tau \circ \Delta(a)\mathcal{R}, \quad a \in U_q(\hat{sl}(2, C)), \quad (4.3)$$

where τ is the permutation operator $\tau(t_1 \otimes t_2) = t_2 \otimes t_1$, for $t_1, t_2 \in U_q(\hat{sl}(2, C))$. For simplicity we sometimes write $U_q(\hat{sl}(2, \mathbf{C}))$ and $U_q(sl(2, \mathbf{C}))$ by $\hat{U}_q$ and U_q , respectively.

Let us discuss solvable models from the viewpoint of $\hat{U}_q$. We define a homomorphism $\phi: \hat{U}_q \rightarrow U_q$ by $\phi(X_0^\pm) = X^\mp$, $\phi(X_1^\pm) = X^\pm$, and $\phi(H_0) = -H$, $\phi(H_1) = H$. We introduce an automorphism $T_x: \hat{U}_q \rightarrow \hat{U}_q$ by $T_x(X_0^\pm) = x^{\pm 1} X_0^\pm$, $T_x(X) = X$, for $X = X_1^\pm, H_0, H_1$.

We define an operator $R(x)$ by $R(x) = \phi((T_x \otimes I)(\mathcal{R}))$. Then the operator $R(x)$ satisfies the following for $i = 0, 1$.

$$R(x)q^{\hat{H}_i/2} \otimes q^{\hat{H}_i/2} = q^{\hat{H}_i/2} \otimes q^{\hat{H}_i/2} R(x), \quad (4.4)$$

$$\begin{aligned}
& R(x)(x^{\delta(i,0)} \hat{X}_i^+ \otimes q^{\hat{H}_i/2} + q^{-\hat{H}_i/2} \otimes \hat{X}_i^+) \\
&= (x^{\delta(i,0)} \hat{X}_i^+ \otimes q^{-\hat{H}_i/2} + q^{\hat{H}_i/2} \otimes \hat{X}_i^+) R(x), \tag{4.5}
\end{aligned}$$

$$\begin{aligned}
& R(x)(x^{-\delta(i,0)} \hat{X}_i^- \otimes q^{\hat{H}_i/2} + q^{-\hat{H}_i/2} \otimes \hat{X}_i^-) \\
&= (x^{-\delta(i,0)} \hat{X}_i^- \otimes q^{-\hat{H}_i/2} + q^{\hat{H}_i/2} \otimes \hat{X}_i^-) R(x). \tag{4.6}
\end{aligned}$$

Here $\hat{X}_i^\pm = \phi(X_i^\pm)$, $\hat{H}_i = \phi(H_i)$ for $i = 0, 1$.

Let us take arbitrary two representations μ_1, μ_2 of U_q . We can take as μ_1 and μ_2 the spin j representations of $U_q(sl(2))$, or the (infinite dim. or finite dim.) color representations of $U_q(sl(2))$. We define R matrix and its matrix elements for the representations by

$$\begin{aligned}
R_{\mu_1 \mu_2}(x) &= \pi^{\mu_1} \otimes \pi^{\mu_2}(R(x)), \\
R_{\mu_1 \mu_2}(x)_{b_1 b_2}^{a_1 a_2} &= (\pi^{\mu_1} \otimes \pi^{\mu_2}(R(x)))_{b_1 b_2}^{a_1 a_2}. \tag{4.7}
\end{aligned}$$

Then the relations (4.4), (4.5) and (4.6) give linear equations for the R matrix elements. It is easy to see that if the matrix elements satisfy the linear equations, then the matrix elements satisfy the Yang-Baxter equations. [27]

$$\begin{aligned}
& \sum_{c_1 c_2 c_3} R_{p_1 p_2}(u)_{b_1 b_2}^{c_1 c_2} R_{p_1 p_3}(u+v)_{c_1 b_3}^{a_1 c_3} R_{p_2 p_3}(v)_{c_2 c_3}^{a_2 a_3} \\
&= \sum_{c_1 c_2 c_3} R_{p_1 p_2}(u)_{c_1 c_2}^{a_1 a_2} R_{p_1 p_3}(u+v)_{b_1 c_3}^{c_1 a_3} R_{p_2 p_3}(v)_{b_2 b_3}^{c_2 c_3}. \tag{4.8}
\end{aligned}$$

Here we have defined the spectral parameter u by $x = \exp u$. In operator formalism, solutions of the linear equations (R matrix) can be written as

$$R(q; u) = \sum_{\mu} |\mu_1 \mu_2; \mu\rangle g(\mu_1, \mu_2, \mu; u) \langle \mu_1 \mu_2; \mu| \tag{4.9}$$

Here the sum is taken over all μ appearing in the decomposition of the tensor product $\mu_1 \otimes \mu_2$ ($\mu \subset \mu_1 \otimes \mu_2$), and $g(\mu_1, \mu_2, \mu; u)$ is some function of the spectral parameter $x = \exp u$.

4.2 Infinite dimensional case

Let us introduce the colored vertex models. [3, 6] The Boltzmann weights of the colored vertex model $X_{\alpha\beta}(u)_{cd}^{ab}$ is defined for the configuration $\{a, b, c, d\}$.

(Fig. 1) The Yang-Baxter relation for the colored vertex model reads

$$\begin{aligned} & \sum_{c_1, c_2, c_3} X_{\alpha_1, \alpha_2}(u_1 - u_2)_{c_2 c_1}^{a_1 a_2} X_{\alpha_1, \alpha_3}(u_1 - u_3)_{c_3 b_1}^{c_1 a_3} X_{\alpha_2, \alpha_3}(u_2 - u_3)_{b_3 b_2}^{c_2 c_3} \\ &= \sum_{c_1, c_2, c_3} X_{\alpha_2, \alpha_3}(u_2 - u_3)_{c_3 c_2}^{a_2 a_3} X_{\alpha_1, \alpha_3}(u_1 - u_3)_{b_3 c_1}^{a_1 c_3} X_{\alpha_1, \alpha_2}(u_1 - u_2)_{b_2 b_1}^{c_1 c_2}. \end{aligned} \quad (4.10)$$

We construct the colored vertex models by calculating the matrix elements of the R matrix (4.9) on the color representations. Let us consider color representations $(\pi_q^{(p_i)}, V^{(p)})$ ($i = 1, 2$) of $U_q(sl(2))$. We define $R_{p_1 p_2}(q; u)$ and its matrix elements by

$$\begin{aligned} R_{p_1 p_2}(q; u) &= \pi_q^{(p_1)} \otimes \pi_q^{(p_2)}(R(x)), \\ R_{p_1 p_2}(q; u)_{b_1 b_2}^{a_1 a_2} &= \left(\pi_q^{(p_1)} \otimes \pi_q^{(p_2)}(R(x)) \right)_{b_1 b_2}^{a_1 a_2}. \end{aligned} \quad (4.11)$$

Here $a_1, a_2, b_1, b_2 = 0, 1, \dots, \infty$.

We can show that the following gives a solution of the linear equations (4.4), (4.5) and (4.6).

$$\begin{aligned} R_{p_1 p_2}(q; u)_{b_1 b_2}^{a_1 a_2} &= \sum_n^{\infty} g(p_1, p_2, n; u)_q \\ &\quad C(p_1, p_2, p_1 + p_2 - n; b_1, b_2, b_1 + b_2 - n)_q \\ &\quad C(p_2, p_1, p_1 + p_2 - n; a_2, a_1, a_1 + a_2 - n)_q. \end{aligned} \quad (4.12)$$

where

$$g(p_1, p_2, n; u)_q = (-1)^n \prod_{k=0}^{n-1} \frac{[u - p_1 - p_2 + k]_q}{[u + p_1 + p_2 - k]_q}, \text{ for } n \in \mathbf{Z}_{\geq 0}. \quad (4.13)$$

We assume $g(p_1, p_2, n = 0; u)_q = 1$. For the fixed values of $a_i, b_i (i = 1, 2)$, the sum in (4.12) reduces to a finite sum. Therefore the sum in (4.12) is well defined. Thus we have an explicit formula for the R matrix elements for the infinite dimensional color representation. It is remarked that the case $p_1 = p_2$, the infinite dimensional representation of the R matrix with a spectral parameter were discussed by Jimbo by using projection operators. [22]

Through the R matrix (4.12) the Boltzmann weights of the colored vertex model are written as follows

$$X_{\alpha\beta}^{(\infty)}(u)_{cd}^{ab} = R_{p_1 p_2}(q; u)_{cd}^{ba}, \quad (4.14)$$

where $\alpha = q^{p_2}$ and $\beta = q^{p_1}$. Thus we obtain the colored vertex model of the infinite state case introduced in the reference [6].

Using the Clebsch-Gordan coefficients we can calculate R matrix for the representations V^j and $V^{(\infty)}$. We define $R_{p_1 j_2}(x)$ and its matrix elements by

$$\begin{aligned} R_{p_1 j_2}(x) &= \pi_q^{(p_1)} \otimes \pi_q^{(j_2)}(R(x)), \\ R_{p_1 j_2}(x)_{b_1 b_2}^{a_1 a_2} &= \left(\pi_q^{(p_1)} \otimes \pi_q^{(j_2)}(R(x)) \right)_{cd}^{ab}. \end{aligned} \quad (4.15)$$

Here $a_1, b_1 = 0, 1, \dots, N-1$, and $a_2, b_2 = 0, 1, \dots, 2j_2$. The matrix elements are given by the following.

$$\begin{aligned} R_{p_1 j_2}(x)_{b_1 b_2}^{a_1 a_2} &= \sum_{n=0}^{2j_2} g(p_1, j_2, n; u)_q \\ &\quad C(p_1, j_2, p_1 + p_2 - n; b_1, b_2, b_1 + b_2 - n)_q \\ &\quad C(j_2, p_1, p_1 + p_2 - n; a_2, a_1, a_1 + a_2 - n)_q. \end{aligned} \quad (4.16)$$

Here $g(p_1, j_2, n; u)_q$ is given by (4.13) with $p_2 = j_2$.

4.3 Finite dimensional case

Let us consider color representations $(\pi_\epsilon^{(p_i)}, V^{(N)})$ ($i = 1, 2$) of $U_q(sl(2))$ with $q = \epsilon$. We define $R_{p_1 p_2}(\epsilon; x)$ and its matrix elements by

$$\begin{aligned} R_{p_1 p_2}(\epsilon; x) &= \pi_\epsilon^{(p_1)} \otimes \pi_\epsilon^{(p_2)}(R(x)), \\ R_{p_1 p_2}(\epsilon; x)_{b_1 b_2}^{a_1 a_2} &= \left(\pi_\epsilon^{(p_1)} \otimes \pi_\epsilon^{(p_2)}(R(x)) \right)_{b_1 b_2}^{a_1 a_2}. \end{aligned} \quad (4.17)$$

Then we have the following.

$$\begin{aligned} R_{p_1 p_2}(\epsilon; u)_{b_1 b_2}^{a_1 a_2} &= \sum_{n=0}^{N-1} g(p_1, p_2, n; u)_\epsilon \\ &\quad \times C(p_1, p_2, p_1 + p_2 - n; b_1, b_2, b_1 + b_2 - n)_\epsilon \\ &\quad \times C(p_2, p_1, p_1 + p_2 - n; a_2, a_1, a_1 + a_2 - n)_\epsilon, \end{aligned} \quad (4.18)$$

where

$$g(p_1, p_2, n; u)_\epsilon = (-1)^n \prod_{k=0}^{n-1} \frac{[u - p_1 - p_2 + k]_\epsilon}{[u + p_1 + p_2 - k]_\epsilon}, \text{ for } n \in \mathbf{Z}_{\geq 0}. \quad (4.19)$$

We assume $g(p_1, p_2, n = 0; u)_\epsilon = 1$.

Through the R matrix (4.12) the Boltzmann weights of the colored vertex model are written as follows

$$X_{\alpha\beta}^{(N)}(u)_{cd}^{ab} = R_{p_1 p_2}(\epsilon; u)_{cd}^{ba}, \quad (4.20)$$

where $\alpha = \epsilon^{p_2}$ and $\beta = \epsilon^{p_1}$. Thus we obtain the colored vertex model of the finite state case given in the reference [3].

Using the Clebsch-Gordan coefficients we can calculate R matrix for the representations V^j and $V^{(N)}$. We define $R_{p_1 j_2}(x)$ and its matrix elements by

$$\begin{aligned} R_{p_1 j_2}(\epsilon; x) &= \pi_\epsilon^{(p_1)} \otimes \pi_\epsilon^{(j_2)}(R(x)), \\ R_{p_1 j_2}(\epsilon; x)_{b_1 b_2}^{a_1 a_2} &= \left(\pi_\epsilon^{(p_1)} \otimes \pi_\epsilon^{(j_2)}(R(x)) \right)_{b_1 b_2}^{a_1 a_2}. \end{aligned} \quad (4.21)$$

Here $a_1, b_1 = 0, 1, \dots, N-1$, and $a_2, b_2 = 0, 1, \dots, 2j_2$. The matrix elements are given by the following.

$$\begin{aligned}
R_{p_1 j_2}(\epsilon; x)_{b_1 b_2}^{a_1 a_2} &= \sum_n^{2j_2} g(p_1, j_2, n; u)_\epsilon \\
&\quad C(p_1, j_2, p_1 + p_2 - n; b_1, b_2, b_1 + b_2 - n)_\epsilon \\
&\quad C(j_2, p_1, p_1 + p_2 - n; a_2, a_1, a_1 + a_2 - n)_\epsilon. \quad (4.22)
\end{aligned}$$

Here the function $g(p_1, j_2, n; u)_\epsilon$ is given by (4.19) with $p_2 = j_2$. It is easy to show that the R matrix (4.22) gives a solution of the linear equation (4.5).

5 Colored IRF models

5.1 Infinite dimensional case

Using the Racah coefficients for color representations we define colored IRF (Interaction Round a Face) models. Let the symbol $w(a, b, c, d; p_1, p_2; u)_q$ denotes the Boltzmann weight of colored IRF model for the configuration $\{a, b, c, d; p_1, p_2\}$. (Fig. 2) The admissible condition is determined by the fusion rules. In the configuration $\{a, b, c, d; p_1, p_2\}$, a is admissible to d if $a \subset p_1 \otimes d$, i.e., $N_{p_1, d}^a \neq 0$. We denote this condition by $a \sim d$. The Boltzmann weight $w(a, b, c, d; p_1, p_2; u)_q$ is defined to be zero, unless $a \sim d$, $b \sim a$, $c \sim d$ and $b \sim c$.

The Yang-Baxter relation for the colored IRF model is given by the following.

$$\begin{aligned}
&\sum_g w(g_1, g_2, g, g_0; p_1, p_2; u) w(g_2, g_3, g'_2, g; p_1, p_3; u + v) \\
&\quad w(g, g'_2, g'_1, g_0; p_2, p_3; v) \\
&= \sum_g w(g_2, g_3, g, g_1; p_2, p_3; v) w(g_1, g, g'_1, g_0; p_1, p_3; u + v)
\end{aligned}$$

$$w(g, g_3, g_2', g_1'; p_1, p_2; u). \quad (5.1)$$

The Boltzmann weights of the colored IRF models are given as follows.

$$\begin{aligned} & w(g_{12}, g_1, g_{21}, g_0; p_1, p_2; u)_q \\ &= \sum_{n=0}^{\infty} g(p_1, p_2, n; u)_q [2p+1]_q \sqrt{[2g_{12}+1]_q [2g_{21}+1]_q} \\ & \quad W(p_2, p_1, g_1, g_0; p, g_{12})_q W(p_2, g_1, p_1, g_0; g_{21}, p)_q, \end{aligned} \quad (5.2)$$

where $p = p_1 + p_2 - n, n \in \mathbf{Z}_{\geq 0}$, and the function $g(p_1, p_2, n; u)_q$ is given by (4.13). We recall that the symbols W and w denote the Racah coefficient and the Boltzmann weight of the IRF model, respectively.

The expression (5.2) can be derived from (4.12) by using the definition of the Racah coefficients [28] (see also [29] for the cases of the spin j representations). Therefore the Boltzmann weights given in (5.2) satisfy the Yang-Baxter relation for the IRF models (5.1).

We can discuss hybrid type colored IRF models, i.e., $2\mu_1 \in \mathbf{Z}_{\geq 0}$ or $2\mu_2 \in \mathbf{Z}_{\geq 0}$. For simplicity we consider only the case $p_1 \in \mathbf{C}$ and $2\mu_2 = 2j_2 \in \mathbf{Z}_{\geq 0}$. The Boltzmann weights of the colored IRF models are given as follows.

$$\begin{aligned} & w(g_{12}, g_1, g_{21}, g_0; p_1, j_2; u)_q \\ &= \sum_{n=0}^{2j_2} g(p_1, j_2, n; u)_q [2p+1]_q \sqrt{[2g_{12}+1]_q [2g_{21}+1]_q} \\ & \quad W(j_2, p_1, g_1, g_0; p, g_{12})_q W(p_1, j_2, g_1, g_0; g_{21}, p)_q, \end{aligned} \quad (5.3)$$

where $p = p_1 + j_2 - n, 0 \leq n \leq 2j_2, n \in \mathbf{Z}_{\geq 0}$, and the function $g(p_1, p_2, n; u)_q$ is given by (4.13).

5.2 Finite dimensional case

Using the Racah coefficients for color representations we define colored IRF (Interaction Round a Face) models. Let the symbol $w(a, b, c, d; p_1, p_2; u)_\epsilon$

denote the Boltzmann weight of colored IRF model associated with finite dimensional color representations of $U_q(sl(2))$.

The Boltzmann weights of the colored IRF model are given by the following.

$$\begin{aligned}
& w(g_{12}, g_1, g_{21}, g_0; p_1, p_2; u)_\epsilon \\
&= \sum_{n=0}^{N-1} g(p_1, p_2, n; u)_\epsilon [2p+1]_q \sqrt{[2g_{12}+1]_\epsilon [2g_{21}+1]_\epsilon} \\
& \quad \times W(p_2, p_1, g_1, g_0; p, g_{12})_\epsilon W(p_1, p_2, g_1, g_0; p, g_{21})_\epsilon
\end{aligned} \tag{5.4}$$

Here the function $g(p_1, p_2, n; u)_\epsilon$ is given by (4.19).

We can discuss hybrid type colored IRF models, i.e., $2\mu_1 \in \mathbf{Z}_{\geq 0}$ or $2\mu_2 \in \mathbf{Z}_{\geq 0}$. For simplicity we consider only the case $p_1 \in \mathbf{C}$ and $2\mu_2 = 2j_2 \in \mathbf{Z}_{\geq 0}$. The Boltzmann weights of the colored IRF models are given as follows.

$$\begin{aligned}
& w(g_{12}, g_1, g_{21}, g_0; p_1, j_2; u)_\epsilon \\
&= \sum_{n=0}^{2j_2} g(p_1, j_2, n; u)_\epsilon [2p+1]_\epsilon \sqrt{[2g_{12}+1]_\epsilon [2g_{21}+1]_\epsilon} \\
& \quad W(j_2, p_1, g_1, g_0; p, g_{12})_\epsilon W(p_1, j_2, g_1, g_0; p, g_{21})_\epsilon
\end{aligned} \tag{5.5}$$

where $p = p_1 + j_2 - n$, $0 \leq n \leq 2j_2$, $n \in \mathbf{Z}_{\geq 0}$, and the function $g(p_1, j_2, n)_\epsilon$ is given by (4.19).

6 Invariants of trivalent colored oriented graphs

6.1 Basic relations for the colored braid matrices

We construct new invariants of colored oriented graphs using the Clebsch-Gordan coefficients for finite dimensional color representations given in §3.

Let us define the N -state colored braid matrix. We assume the charge conservation: $G_{cd}^{ab}(p_\alpha, p_\beta; \pm) = 0$ unless $a + b = c + d$ ($\alpha = q^{-4p_\alpha}$, $\beta = q^{-4p_\beta}$).

Furthermore we assume $G_{cd}^{ab}(p_\alpha, p_\beta; +) = 0$ if $a < d$, and $G_{cd}^{ab}(p_\alpha, p_\beta; -) = 0$ if $a > d$. We define $(z; q)_n$ by $(z; q)_n = (1 - z)(1 - zq) \cdots (1 - zq^{n-1})$ for $n > 0$ and $(z; q)_0 = 1$ for $n = 0$. The nonzero elements are written as follows.

$$\begin{aligned}
G_{c,d}^{a,b}(p_\alpha, p_\beta; +) &= \epsilon^{-2bp_\alpha - 2dp_\beta} \epsilon^{2bd} \epsilon^{2p_\alpha p_\beta} \\
&\left(\frac{(\epsilon^2; \epsilon^2)_c}{(\epsilon^2; \epsilon^2)_b (\epsilon^2; \epsilon^2)_{c-b}} \frac{(\epsilon^2; \epsilon^2)_a}{(\epsilon^2; \epsilon^2)_d (\epsilon^2; \epsilon^2)_{a-d}} \frac{(\epsilon^{-4p_\alpha}; \epsilon^2)_a}{(\epsilon^{-4p_\alpha}; \epsilon^{-2})_d} \frac{(\epsilon^{-4p_\beta}; \epsilon^2)_c}{(\epsilon^{-4p_\beta}; \epsilon^2)_b} \right)^{1/2}, \\
G_{c,d}^{a,b}(p_\alpha, p_\beta; -) &= \epsilon^{2ap_\alpha + 2cp_\beta} \epsilon^{-2ac} \epsilon^{-2p_\alpha p_\beta} \\
&\left(\frac{(\epsilon^{-2}; \epsilon^{-2})_b}{(\epsilon^{-2}; \epsilon^{-2})_{b-c} (\epsilon^{-2}; \epsilon^{-2})_c} \frac{(\epsilon^{-2}; \epsilon^{-2})_d}{(\epsilon^{-2}; \epsilon^{-2})_{d-a} (\epsilon^{-2}; \epsilon^{-2})_a} \frac{(\epsilon^{4p_\alpha}; \epsilon^{-2})_b}{(\epsilon^{4p_\alpha}; \epsilon^{-2})_c} \frac{(\epsilon^{4p_\beta}; \epsilon^{-2})_d}{(\epsilon^{4p_\beta}; \epsilon^{-2})_a} \right)^{1/2}.
\end{aligned} \tag{6.1}$$

Then the colored braid matrix satisfies the following relation.

$$\begin{aligned}
&\sum_{c_1, c_2, c_3} G_{c_2, c_1}^{a_1, a_2}(p_1, p_2; +) G_{c_3, b_1}^{c_1, a_3}(p_1, p_3; +) G_{b_3, b_2}^{c_2, c_3}(p_2, p_3; +) \\
&= \sum_{c_1, c_2, c_3} G_{c_3, c_2}^{a_2, a_3}(p_2, p_3; +) G_{b_3, c_1}^{a_1, c_3}(p_1, p_3; +) G_{b_2, b_1}^{c_1, c_2}(p_1, p_2; +).
\end{aligned} \tag{6.2}$$

Here p_1, p_2, p_3 denote the colors of the strings.

It has been explicitly shown that the colored braid matrix is equivalent to the R matrix of the color representations of $U_q(sl(2))$. [6]

Let us introduce basic relations for the colored braid matrix and the Clebsch-Gordan coefficients. We note they are related to the Reidemeister moves.

(i) First inversion relation:

$$\sum_{e,f} G_{ef}^{ab}(p_1, p_2; +) G_{cd}^{ef}(p_1, p_2; -) = \sum_{e,f} G_{ef}^{ab}(p_1, p_2; -) G_{cd}^{ef}(p_1, p_2; +) = \delta_c^a \delta_d^b. \tag{6.3}$$

(ii) Second inversion relation:

$$\sum_{e,f} G_{cf}^{ae}(p_1, p_2; +) G_{be}^{df}(p_1, p_2; -) \omega^{d-e} = \delta_b^a \delta_d^c,$$

$$\sum_{e,f} G_{cf}^{ae}(p_2, p_1; -) G_{be}^{df}(p_2, p_1; +) \omega^{d-e} = \delta_b^a \delta_d^c. \quad (6.4)$$

(iii) Markov trace property:

$$\sum_b G_{ab}^{ab}(p, p; \pm) \omega^{-b} = \epsilon^{2p(N-1) \pm 2p(N-1) \pm 2p^2}. \quad (6.5)$$

(iv)

$$\begin{aligned} & \sum_{c_1 c_2} G_{b_3 c_1}^{a_1 c_3}(p_1, p_3; +) G_{c_3 c_2}^{a_2 a_3}(p_2, p_3; +) C(p_1, p_2, p; c_1, c_2, b)_\epsilon \\ &= \sum_c C(p_1, p_2, p; a_1, a_2, c)_\epsilon G_{b_3 b}^{ca_3}(p, p_3; +), \\ & \sum_{c_1 c_2} G_{b_3 c_1}^{a_1 c_3}(p_3, p_1; -) G_{c_3 c_2}^{a_2 a_3}(p_3, p_2; -) C(p_1, p_2, p; c_1, c_2, b)_\epsilon \\ &= \sum_c C(p_1, p_2, p; a_1, a_2, c)_\epsilon G_{b_3 b}^{ca_3}(p_3, p; -). \end{aligned} \quad (6.6)$$

6.2 Invariants of colored oriented framed tangle graphs with trivalent vertices

We consider trivalent graphs, which have vertices with three edges. Framed graphs have framing vector fields. We assume that all three edges have common tangent vector in the trivalent vertex, and the framing vector field is always normal to the tangent vector to the vertex. (Fig. 3)

We introduce "tangle graph". We define (k, l) -oriented tangle graph T by a finite set of disjoint oriented arcs, oriented trivalent vertices, and oriented circles properly embedded in $\mathbf{R}^2 \times [0, 1]$ such that

$$\partial T = \{(i, 0, 0); i = 1, 2, \dots, k\} \cup \{(j, 0, 1); j = 1, \dots, l\}, \quad (6.7)$$

where ∂T denotes the upper and lower boundaries of the tangle graph. (Fig. 4) We define colored oriented tangle graphs (T, α) by assigning colors on edges, arcs and circles of tangle graphs.

We can express trivalent framed tangle graph by plane diagram. [30, 31, 32] We can choose the diagram such that the framing vector field is normal to the plane of projection and directed "up".

Proposition 6.1 [30] *The isotopy invariants of framed tangle graphs are functions on their plane diagrams, invariant under the following local moves $D_1 \sim D_7$.*

(Fig. 5)

We introduce weights for diagrams. We assign the colored braid matrices, the Clebsch-Gordan coefficients, and the weights U_i to the braiding diagrams, trivalent vertex diagrams, and the creation-annihilation diagrams, respectively. (Fig. 6)

We define $\phi(T, \alpha)$ for a tangle graph T by the summation over all possible configurations of variables z_i on the edges (or segments) of the graph. In the summation we fix the colors p_i . The sum corresponds to partition function in statistical mechanics. Then from the relations (6.2), (6.3), (6.4), (6.5) the sum $\phi(T, \alpha)$ for the tangle graph T is invariant under the moves $D_1 \sim D_7$. Thus we have isotopy invariants of trivalent colored oriented framed tangle graphs.

6.3 Invariants of trivalent framed graphs

We construct invariants of trivalent colored oriented framed graphs by an approach parallel to that for the colored link invariants. [5] Through $(1, 1)$ -tangle graph we introduce another invariant of a framed graph. [5]

Let T be a $(1, 1)$ -tangle graph. We denote by $\hat{T}$ the graph obtained by closing the open strings of T . It is easy to show the following proposition. [5]

Proposition 6.2 *Let T_1 and T_2 denote two $(1, 1)$ -tangle graph. If $\hat{T}_1$ is isotopic to $\hat{T}_2$ as a graph in S^3 by an isotopy which carries the closing component of $\hat{T}_1$ to that of $\hat{T}_2$. Then T_1 is isotopic to T_2 as a $(1, 1)$ -tangle graph.*

Let T be a $(1, 1)$ -tangle graph. We put $F = \hat{T}$ and s is the color of the closing component (or edge) of $\hat{T}$. We denote by $\phi(T, \alpha)_b^a$ the value ϕ for the tangle graph with variables a and b on the closing component (or edge). (Fig. 7)

Then from the discussion given in the reference [5] we can show that

$$\phi(T, \alpha)_b^a = \lambda \delta_{ab}. \quad (6.8)$$

The value of $\phi(T, \alpha)_b^a$ do not depend on a or b .

For a colored graph (F, α) and a color s of closing component (or edge), we define Φ by $\Phi(F, s, \alpha) = \lambda$ where F, T, s are above and $\phi(T, \alpha)_b^a = \lambda \delta_{ab}$. By the basic relations given in §6.2, Φ is well-defined, i.e. $\Phi(F, s, \alpha)$ does not depend on a choice of T .

Further we have the next proposition, to obtain invariants which do not depend on s .

Proposition 6.3 [5] *For a graph F and its color $\alpha = (p_1, \dots, p_n)$, we have the next formula.*

$$\Phi(F, s, \alpha)([p_s; N-1]_\epsilon!)^{-1} = \Phi(F, s', \alpha)([p_{s'}; N-1]_\epsilon!)^{-1}. \quad (6.9)$$

The proof of this proposition is equivalent to that given in Appendix C of the reference [5]. By this proposition we obtain the next definition.

Definition 6.4 *For a trivalent colored oriented framed graph (F, α) , we define an isotopy invariant $\hat{\Phi}$ of (F, α) by*

$$\hat{\Phi}(F, \alpha) = \Phi(F, s, \alpha)([p_s; N-1]_\epsilon!)^{-1}. \quad (6.10)$$

Thus we obtain new invariants $\Phi(F, \boldsymbol{\alpha})$ of trivalent colored oriented framed graphs $(F, \boldsymbol{\alpha})$.

Acknowledgement

We would like to thank Prof. M. Wadati for encouragement.

A Appendix A

We give derivation of the Clebsch-Gordan coefficients for infinite dimensional representations of $U_q(sl(2))$. We calculate the Clebsch-Gordan coefficients following Racah's approach [20]. For simplicity we denote $C(p_1, p_2, p_1 + p_2 - n; z_1, z_2, z)_q$ by $C(n; z_1, z_2, z)$.

Applying $\Delta(X^\pm)$ on $|p_1, p_2; p, z\rangle$ ($p = p_1 + p_2 - n$), we have

$$\begin{aligned} & \sqrt{[z][2p+1-z]}C(n; z_1, z_2-1, z-1) \\ = & \sqrt{[z_1+1][2p_1-z_1]}q^{p_2-z_2+1}C(n; z_1+1, z_2-1, z) \\ & + q^{-(p_1-z_1)}\sqrt{[z_2][2p_2+1-z_2]}C(n; z_1, z_2, z), \end{aligned} \quad (A.1)$$

$$\begin{aligned} & \sqrt{[z+1][2p-z]}C(n; z_1, z_2+1, z+1) \\ = & \sqrt{[z_1][2p_1+1-z_1]}q^{p_2-z_2-1}C(n; z_1-1, z_2+1, z) \\ & + q^{-(p_1-z_1)}\sqrt{[z_2+1][2p_2-z_2]}C(n; z_1, z_2, z). \end{aligned} \quad (A.2)$$

We introduce $f(n; z_1, z_2, z)$ by

$$\begin{aligned} C(n; z_1, z_2, z) &= (-1)^{z_1}q^{(p_1-z_1)(p-z+1)} \\ &\times \left(\frac{[2p_1-z_1; n-z_1]![2p_2-z_2; n-z_2]!}{[2p; z]![z_1]![z_2]![z]!} \right)^{1/2} f(n; z_1, z_2, z). \end{aligned} \quad (A.3)$$

Then we have the following recurrence relations.

$$\begin{aligned} f(n; z_1, z_2+1) &= -[z_1][2p_1+1-z_1]q^{2p-2z}f(n; z_1-1, z_2+1, z) \\ &+ [2p_2-z_2][z_2+1]f(n; z_1, z_2), \end{aligned} \quad (A.4)$$

$$\begin{aligned} & [z][2p-z+1]q^{2(p_1-z_1)}f(n; z_1, z_2-1) \\ = & -f(n; z_1+1, z_2-1, z) + f(n; z_1, z_2, z). \end{aligned} \quad (A.5)$$

Setting $z = 0$ in (A.5) we have $f(n; z_1, z_2, 0) = f(n; z_1+1, z_2-1, 0)$. Therefore we write $f(n; z_1, z_2, 0)$ as f_n . From (A.4) we have

$$f(n; z_1, z_2, z) = f_n[z_1]![z_2]!\sum_t (-1)^t q^{t(2p+1-z)} \begin{bmatrix} z \\ t \end{bmatrix}$$

$$\times \frac{[2p_1 - z_1 + t; t]![2p_2 - z_2 + z - t; z - t]!}{[z_1 - t]![z_2 - z + t]!}. \quad (\text{A.6})$$

By using (B.3) it is easy to show that (A.6) satisfies the recurrence relation (A.4).

Now we consider f_n . If we assume the following expression of $C(n; z_1, z_2, 0)$

$$C(n; z_1, z_2, 0) = (-1)^{z_1} q^{n(1-n)/2 - z_1(p+1) + np_2} \times \left(\frac{[n]![2p_1 - z_1; n - z_1]![2p_2 - z_2; n - z_2]!}{[2p + n + 1; n]![z_1]![z_2]!} \right)^{1/2}, \quad (\text{A.7})$$

we can show the orthogonality relation

$$\sum_{z_1} C(m; z_1, n - z_1, n - m) C(n; z_1, n - z_1, 0) = \delta_{nm}, \quad 0 \leq m \leq n. \quad (\text{A.8})$$

We prove (A.8) in the following. When $m = n$, we show (A.8) using (B.6) where $\alpha = 2p_1 - n$ and $\beta = 2p_2 - n$. When $m < n$, we prove (A.8) putting the expression (A.6) and (A.7) in the relation (A.8), transforming the variables $\{z_1, t\}$ into $\{t, \nu\}$ ($z_1 - t = \nu$) and then taking the sum over t using (B.4). Then we see that the sum vanishes.

From the relation (A.8) and the convention of phase factor, the expression (A.7) is shown. We obtain f_n as follows.

$$f_n = q^{n(1-n)/2 + np_2 - p_1(p+1)} \sqrt{\frac{[n]!}{[2p + n + 1; n]!}}. \quad (\text{A.9})$$

Thus we have the following expression for the Clebsch-Gordan coefficients.

$$\begin{aligned} C(n; z_1, z_2, z) &= \delta(z_1 + z_2 - n, z) q^{n(1-n)/2 + np_2 - z_1(p+1) - (p_1 - z_1)z} \\ &\times \left(\frac{[2p_1 - z_1; n - z_1]![2p_2 - z_2; n - z_2]![n]![z_1]![z_2]![z]!}{[2p + n + 1; n]![2p; z]!} \right)^{1/2} \\ &\sum_t (-1)^{t+z_1} q^{t(2p+1-z)} \frac{[2p_1 - z_1 + t; t]![2p_2 - z_2 + z - t; z - t]!}{[t]![z - t]![z_1 - t]![z_2 - z + t]!}. \end{aligned} \quad (\text{A.10})$$

Finally we transform the expression (A.10) of the Clebsch-Gordan coefficients into the symmetric one (3.10) given in the section 3. We apply the formula (B.5) to the expression (A.10) ($\alpha = 2p_1 - z_1, \beta = t, c = n - z_1 + t$), and replace the variable k by $k = t - u$. We apply the formula (B.7) ($\alpha = 2p_2 + z - z_2 - u, s = t - u, a = z_1 - u, b = z - u$). Then we replace the variable u by $\nu = z_1 - u$. We thus obtain the symmetric expression (3.10).

We have considered only the case $p_i \in \mathbf{C}$. The Clebsch-Gordan coefficients for the other cases such as $(\mu_1, \mu_2) = (p_1, j_2)$ can be derived in the same way.

B Appendix B

We define q -analog of combinatorial for $m \geq k, m, k \in \mathbf{Z}_{\geq 0}$ and $\alpha \in \mathbf{C}$ as follows.

$$\begin{bmatrix} m \\ k \end{bmatrix} = \frac{[m]!}{[m-k]![k]!}, \quad \begin{bmatrix} \alpha \\ k \end{bmatrix} = \frac{[\alpha; k]!}{[k]!}. \quad (\text{B.1})$$

(1) For $m, n \in \mathbf{Z}_{\geq 0}$ ($n \geq m$), we have the following useful formulas.

$$\prod_{k=1}^m (1 - zq^{2k-2}) = \sum_k (-z)^k q^{k(m-1)} \begin{bmatrix} m \\ k \end{bmatrix}, \quad (\text{B.2})$$

$$\begin{bmatrix} n-1 \\ m \end{bmatrix} + \begin{bmatrix} n-1 \\ m-1 \end{bmatrix} q^{\pm n} = \begin{bmatrix} n \\ m \end{bmatrix} q^{\pm m}, \quad (\text{B.3})$$

$$\sum_{k=0}^{n-m} (-1)^k \begin{bmatrix} n-m \\ k \end{bmatrix} q^{k(n-m-1)} = \delta_{mn}. \quad (\text{B.4})$$

(2) For $\alpha, \beta \in \mathbf{C}$, and $m, n, a, b, c \in \mathbf{Z}_{\geq 0}$, we have the following.

$$\sum_k \begin{bmatrix} \alpha \\ c-k \end{bmatrix} \begin{bmatrix} \beta \\ k \end{bmatrix} q^{\pm k(\alpha+\beta)} = \begin{bmatrix} \alpha+\beta \\ c \end{bmatrix} q^{\pm c\beta}, \quad (\text{B.5})$$

$$\sum_k \begin{bmatrix} \alpha+n-k \\ n-k \end{bmatrix} \begin{bmatrix} \beta+k \\ k \end{bmatrix} q^{-k(\alpha+\beta+2)} = \begin{bmatrix} \alpha+\beta+n+1 \\ n \end{bmatrix} q^{-n(1+\beta)}, \quad (\text{B.6})$$

$$\sum_s (-1)^s \begin{bmatrix} \alpha - s \\ b - s \end{bmatrix} \begin{bmatrix} a \\ s \end{bmatrix} q^{-s(-\alpha+a+b-1)} = \begin{bmatrix} \alpha - a \\ b \end{bmatrix} q^{-ab}. \quad (\text{B.7})$$

We can derive (B.2) induction on m . We derive the relation (B.3) from (B.2). The relation (B.4) follows (B.2).

We show the relation (B.5) first for the case $\alpha = m_1, \beta = m_2$ ($m_1, m_2 \in \mathbf{Z}_{\geq 0}$) by using (B.2). Since m_1, m_2 are arbitrary positive integers, and (B.5) is equivalent to a polynomial relation in terms of q^α and q^β , the relation (B.5) holds also for complex parameters α, β . From (B.5) we derive the relations (B.6) and (B.7) using the following

$$\begin{bmatrix} \alpha \\ n \end{bmatrix} = (-1)^n \begin{bmatrix} -\alpha - 1 + n \\ n \end{bmatrix}, \quad \text{for } \alpha \in \mathbf{C}, n \in \mathbf{Z}_{\geq 0}. \quad (\text{B.8})$$

References

- [1] Y. Akutsu and T. Deguchi, Phys. Rev. Lett. **67** (1991) 777.
- [2] T. Deguchi and Y. Akutsu, J. Phys. Soc. Jpn. **60** (1991) 2559.
- [3] T. Deguchi and Y. Akutsu, J. Phys. Soc. Jpn. **60** (1991) 4053.
- [4] T. Deguchi, J. Phys. Soc. Jpn. **60** (1991) 3979.
- [5] Y. Akutsu, T. Deguchi and T. Ohtsuki, Invariants of Colored Links, UTYO Math 91-24, preprint 1991 (to appear in Knot Theory and its Ramifications).
- [6] T. Deguchi and Y. Akutsu, Colored braid matrices from infinite dimensional representations of $U_q(g)$, preprint 1991.
- [7] P. Roche and D. Arnaudon, Lett. Math. Phys. **17** (1989) 295-300.
- [8] C. De Concini and V.G. Kac, Prog. Math. **92** (1990) pp. 471-506.
- [9] C. De Concini, V.G. Kac and C. Procesi, Quantum Coadjoint Action, (Preprint di Matematica - n. 95, Scuola Normale Superiore, Pisa), preprint 1991.
- [10] B.U. Felderhof, Physica **66** (1973) 279-297.
- [11] C. Fan and F.Y. Wu: Phys. Rev. **179**(1969) 650; Phys. rev. **B2** (1970) 723.
- [12] T.Deguchi, J. Phys. Soc. Jpn. **58** (1989) 3441-3444;
T.Deguchi and Y. Akutsu, J. Phys. **A: Math. Gen.** **23** (1990) 1861.
- [13] C. L. Schultz: Phys. Rev. Lett. **46** (1981) 629.

- [14] J.H.H. Perk and C.L. Schultz, Phys. Lett. **84A** (1981) 407, and in "Non-Linear Integrable Systems, Classical Theory and Quantum Theory", ed. by M.Jimbo and T.Miwa, World Science, Singapore, 1981.
- [15] P.P. Kulish, E.K. Sklyanin: J. Sov. Math. **19** (1982) 1596.
- [16] L. Rozansky and H. Saleur, Quantum field theory for the multivariable Alexander Conway polynomial, YCTP-P20-91, preprint 1991;
L.H. Kauffman and H. Saleur, Fermions and Link Invariants, YCTP-P21-91, preprint 1991.
- [17] J. Murakami, The Free-Fermion Model in Presence of Field Related to the Quantum Group $U_q(\hat{sl}(2))$ of Affine Type and the Multi-Variable Alexander Polynomial of Links, RIMS-822, preprint 1991.
- [18] J. Murakami, A state model for the multi-variable Alexander polynomial, preprint 1990.
- [19] M. Couture, H.C. Lee, and N.C. Schmeing, in "Physics Geometry and Topology", ed. H.C. Lee, (Plenum Press, New York, 1990), 573-582.
- [20] M.E. Rose, *Elementary Theory of Angular Momentum*, John Wiley and Sons, Inc., New York, 1957.
- [21] V.G.Drinfeld, "Quantum Groups" Proceedings of the ICM, Berkeley (1987) 798 - 820.
- [22] M. Jimbo, Lett. Math. Phys. **10** (1985) 63-69.
- [23] A.N. Kirillov and N.Yu. Reshetikhin, in "Infinite Dimensional Lie Algebras and Groups", ed. V.G. Kac, World Scientific, 1989, p. 285.
- [24] M. Nomura, J. Math. Phys. **30**(1989) 2397-2405.

- [25] N. Yu Reshetikhin and V.G. Turaev, Invent. Math. (1991)
- [26] D. Arnaudon, Phys. Lett. **B268** (1991) 217.
- [27] M. Jimbo, Commun. Math. Phys. **102** (1986) 537-547.
- [28] V. Pasquier, Commun. Math. Phys. **118** (1988) 335.
- [29] M. Nomura, J. Phys. Soc. Jpn. **58** (1989) 2694.
- [30] N. Yu. Reshetikhin and V.G. Turaev, Commun. Math. Phys. **127** (1990) 1-26.
- [31] Louis H. Kauffman, Trans. Amer. Math. Soc. **318** (1990) 317-371.
- [32] Sergey Piunikhin, State sum models for trivalent knotted graph invariants using quantum groups $SL_q(2)$, preprint 1991.

Figure Captions

Fig. 1

(a) Boltzmann weight $X_{\alpha\beta}(u)_{cd}^{ab}$ and (b) R matrix $R_{p_1 p_2}(u)_{b_1 b_2}^{a_1 a_2}$

Fig. 2

Boltzmann weight $w(a, b, c, d; p_1, p_2; u)$

Fig. 3

Trivalent vertex. The edges have colors p_1, p_2, p and variables z_1, z_2, z

Fig. 4

(2,3)-tangle graph. Each segment in the graph has a color p_j .

Fig. 5

Relations $D1 \sim D7$.

Fig. 6

We assign the following weights to the diagrams.

(a) Identity diagram. $I_b^a = \delta_{ab}$, $(I^*)_b^a = \delta_{ab}$

(b) Creation-annihilation diagrams.

$$(U_r)_{ab} = q^{-p(N-1)} \epsilon^{-b} \delta_{a+b, N-1}$$

$$(U_l)_{ab} = q^{p(N-1)} \epsilon^a \delta_{a+b, N-1}$$

$$(\bar{U}_r)_{ab} = q^{p(N-1)} \epsilon^a \delta_{a+b, N-1}$$

$$(\bar{U}_l)_{ab} = q^{-p(N-1)} \epsilon^{-b} \delta_{a+b, N-1}$$

(c) Braiding diagrams

$$G(p_1, p_2; +)_{cd}^{ab} \text{ and } G(p_1, p_2; -)_{cd}^{ab}$$

(d) Vertex diagrams.

We assign the Clebsch-Gordan coefficient $C(p_1, p_2, p; z_1, z_2, z)$. both to the two vertex diagrams V and V' .

Fig. 7

The color s of the closing edge (or component) is p_3 .